

Research article

Phosphogypsum effect on soil, rainwater incubated and soil respiration

Kaouther Ardhaoui^{a,b,*}, Mohamed Moussa^a

^a Institute of arid region, Mednine, Departement of Soil Science, Tunisia, 4100, Mednine

^b University of Gabes, ISBAM, Route El Jorf-km 22.5, Medenine 4119, Tunisia.

* Corresponding author. Tel.: +216 97201985. E-mail address: ardhaouikaouther@gmail.com

Article history:

Received 22 July 2019; Received in revised form 20 August 2019.

Accepted 26 August 2019; Available online 2 September 2019.

Abstract

Phosphogypsum (PG) is a byproduct issued by fertilizer industry, which has been used as soil fertilizer in many countries in the world for long time. The most positive effect of phosphogypsum is ameliorating soil properties and increasing crop yields. Recently, disquiets are growing related to the increase of the acidity and metal ions concentration in soils. This work is a study on the short term impact of using phosphogypsum, as an amendment in El Fja region (Mednine, Tunisia). Physico-chemical characterizations (pH, electrical conductivity, P_2O_5 concentration, chloride concentration, calcium and magnesium concentration) of rain water, incubated at room temperature with soil and also with a (soil, phosphogypsum) mixture during 37 days, were investigated. The dissolution of this product and its impurities in the first period of incubation releases the ions useful for agricultural use like calcium, magnesium, sulphate and phosphate ions. In addition, the effect of amending phosphogypsum on the increase of organic carbon mineralisation, which is an indicator for soil respiration, was revealed.

Key words: Arid zone, phosphogypsum, organic carbon mineralisation, water and soil incubation.

© 2019 Knowledge Journals. All rights reserved.

1. Introduction

Tunisian South Est is an arid zone which is characterized as an area of low precipitation, high temperature, and high rate of evaporation (You, 2016). The soil in this zone is characterized by neutral, high salt content, and low organic matter (Kayouli, 2006; Jeddi, 2010).

Hence, its agronomic potential can be easily weakened by erosion overgrazing or agriculture activity. Subsequently amendments are added to enhance physico-chemical properties of soil (Cherifa, 2009). Several trials have been performed using soil conditioners such as farmyard manure, compost, biochar (Cherifa, 2009; Uchimiya, 2010).

These amendments have beneficial outcome on agricultural productivity, but their effect on soil (microbiological and physico-chemical aspects), water table and global environment are to be investigated seriously. Mineral amendments are known to ameliorate the quality of soil, but their effect is not simultaneous, several weeks are required and even months to notice this effect as reported by the FAO.

This work deals with the use of phosphogypsum PG as an amendment in the arid zone soil by studying its effect on soil-rain water incubated mixtures and soil respiration. In Tunisia, huge amounts of phosphogypsum, estimated to be 10.5 million tons annually (Ben Amor, 2012), are induced by phosphate

fertilizer industry. Hence, numerous trials were performed, with the target of reducing the volumes of PG to be disposed in landfill, and to provide new soil conditioners to balance the use of industrial fertilizers in agriculture especially in poor soil like those in arid zone.

Phosphogypsum dumps constitutes a real environmental struggle as only a small quantity of this production is recycled, although only 15% of world PG production is rehabilitated as building materials (Degirmenci, 2008), as additive in road construction (Sizyakov et al. 2012), as agricultural fertilizers or soil stabilization amendments such as reported by Tayibi (2009). In addition, phosphogypsum can be processed to extract rare-earth metal compounds, because it contains a mass fraction in rare earth oxide Ln_2O_3 varying from 0.1 to 0.8 %, similar to the fraction found in some ores (Akhmetov et al. 1992). Moreover, PG might be treated to produce marketable products like; ammonium sulfate and technical calcium carbonate, calcium sulfide, hydrogen sulfide, sulfuric acid and cement (Genkin et al. 2013).

When it comes to the valorisation of PG in the agriculture, various researches focused their interest on the effect of adding PG into soil. Carvalho, (1997) found that PG can be used as a good amendment for acidic subsoils that present toxic levels of aluminium and/or calcium deficiency. Garrido, (2006) found that

gypsum-rich industrial by-products could regulate the mobility of toxic metals like Cd, Cu and Pb in acid soils. Kumpiene (2008) reported findings about a success in immobilizing toxic pollutant metals for example As, Cr, Cu, Pb and Zn in soil by phosphorus amendments and clays. In a recent study Hentati, (2015) studied the eco-toxicity of amended soil with PG and its impact on bacteria, invertebrates, algae and plants, highlighting the necessity of setting limits for PG application in soils. In a detailed review on the use of Gypsum on Soils, Shainberg (1989) reported the favorable effect of PG on both soil physical properties (especially on dispersive soils), by fighting restricted rooting, and on chemical properties (on acid soils). The PG is also said to be a nutrient source for Ca or S, retainer for water in semiarid or marginal rainfall areas, and a booster to agricultural production.

Otherwise, the richness of soils with organic matter (OM) is one of the main indicators of soil fertility. Soil is an overwhelming living site, so the OM can be involved in the formation or the decomposition processes (Rastetter et al, 1991; Raich et al, 1991). It is also an increasingly feature in a global climate change context where carbon storage in soils is a serious alternative that is predicted to limit CO₂ concentrations in the atmosphere (Wu et al., 2003).

The second part of this research is related to study the kinetic of mineralization, which is an indicator of soil respiration, of the organic matter in an arid zone soil and the impact of adding PG as an amendment on this kinetic. In fact, several researches have been conducted on soil respiration and the effect of adding fertilizers or amendments on this soil activity. Annabi, (2009) showed in a study on the status and dynamics of OM in cultivated and forest soil in the north of Tunisia, that the resistant organic carbon rate to mineralization is bigger in forest soils than in cultivated soils. In another study, Zoghalmi, (2016) found that the amendment of soil with 120 t ha⁻¹ as optimal amount has the best potential for mineralization, which is related to the existing microbial biomass.

Hence, the second part of this work was dedicated to study adding PG and its effects on soil respiration, during 34 days. The kinetic of mineralization of the organic matter is known to be an indicator of the microorganism activity in soil.

2. Materials and methods

El Fje soil has a sandy texture with small amounts of silt (4.92%) and clay (2.17%) which were determined by Robinson pipette (Robinson 1922).

During this study, physico-chemical characterizations (electrical conductivity CE, pH, (Ca²⁺ Mg²⁺) chloride concentrations, and soluble orthophosphate amount) in rainwater were measured. This later was incubated at room temperature with soil and with a (soil, PG: 10% by weight) mixture during 37 days according to the proportions (900 ml rain water, 500g soil with volume ratio 2/1). This

amendment amount 10% of PG in soil was found to be the most suitable for plant development (Ardhaoui, 2017) and closed soil incubation experiments are performed during periods of about one month as stated in literature (Mackay, 1986). The concentration of (Ca²⁺ and Mg²⁺) was determined by Dugain methods (Dugain, 1961).

Organic matter was measured by rapid oxidation and subsequent titration with ferrous ammonium sulphate (Walkley, 1934). Total nitrogen was determined using the Kjeldahl's method (Bremner 1982) by attacking at high temperatures the matter using concentrated sulfuric acid in the presence of a catalyser (MgSO₄, CuSO₄, Se).

The mineralogical components of the PG were identified by X-ray Diffraction (XRD) using a Bruker (D8 advance) powder diffractometer with a tube Cu anode (40 kV, 40 mA).

In an other set of experiments, the amount of carbon dioxide released by soil respiration during 34 days was captured by a (NaOH (1M, 20ml); BaCl₂ (1M, 5ml)) mixture solution maintained during incubation with slightly moisturized soil in a 1 liter glass bottle at 28 °C. The remaining soda concentration, after neutralization with the released CO₂, was obtained by titration with HCl (0.1 M) using phenolphthalein indicator.

3. Results and discussion

Data displayed in a previous research (Ardhaoui et al., 2017) describes the physico-chemical properties of the studied soil, pH is near neutral about 7.5 and the electrical conductivity is (2.33 mS/cm) showing a moderately saline soil (Wong 2009) with a limited cation exchange capacity (4.3 meq/100g), low organic carbon (0.58%) and active carbonate (1.2%) percentages respectively. The total nitrogen amount is 0.18% and that of phosphates (P₂O₅) is 901 ppm.

The Tunisian Chemical Groupe in Gabès furnished Posphogypsum samples. The composition of PG varies depending upon the source of rock phosphate and the process for manufacturing phosphoric acid (Wright 1998). The common composition (Zairi 1999) of PG is: calcium sulfate dihydrate, orthophosphoric acid various salts and trace metals.

It is noteworthy to mention that powdered samples of PG were analyzed by X-ray diffraction, PG is mostly formed by calcium sulfate dihydrate polymorphs and some slight impurities like Mg(OH)₂, SiO₂, MgSiO₃ and Na₂SiF₆ (Figure 1).

Supplementary analysis showed that PG has low pH (2.3) due to the acid remaining (Phosphates P₂O₅ 1209 ppm) from fertilizer manufacture. Additionally, it has an important cation exchange capacity (7.1 meq/100g) and gypsum percentage (97.5%). The organic carbon (1%) and total nitrogen (0.3%) percentages were also obtained.



Fig. 1: X-Ray pattern of powdered PG ($\alpha \text{ Cu}=1.54 \text{ \AA}$)

Fig. 2: Kinetic evolution of physico-chemical properties of incubated rain water with soil and (soil+PG) mixture




Fig. 3: Total amount of carbon dioxide released by soil and (soil+PG) mixture

Furthermore, PG has high ion concentrations due to its mineral composition previously described and to potential crystalline and non-crystalline impurities like chlorate, silicate, phosphate and carbonate known to be present in the phosphate rock generating PG (Drouet 2015; Zairi 1999). All these evidences go with the important EC measured (18 mS/cm).

3.1 Soil incubation in rain water

The Kinetic study (Figure 2) of the physico-chemical properties of rainwater incubated with soil and (soil+ PG) mixture at room temperature showed a significant repercussion of the introduction of PG on these properties. During the period of study, PG amendment enhances EC, (Ca^{2+} , Mg^{2+}) and P_2O_5 concentrations. Chloride ion concentration remained relatively low and the pH decreases while PG is present. This general trend is seen during the first twenty days of incubation. These facts were due, in a first level, to the gradual dissolution of PG producing calcium and sulfate ions and in a second level to the impurities dissolution, that accompany this industrial byproduct. The noteworthy examples of impurities are, iron, aluminum and magnesium sulfates and sulfuric, phosphoric and fluorhydric acids (Zairi, 1999).

During the final period of incubation, the physico-chemical parameters have a tendency to stabilisation and/or the values for the (soil + PG) mixture are getting close to those of soil. At this period, the interaction with soil components begins and then enhances the reactions of precipitation leading to an equilibrium state (Endovitsky, 2016).

These results approve the use of PG as an agriculture fertilizer in different parts of the world for years (Hentati 2015, Mullins 1990). In fact, It has been spread in agricultural soils as a calcium, phosphorus and sulfate supplement to boost crop production and reducing Al-toxicity (Toma 1997) and sodic soils (Enamorado 2014).

It was also used raw or combined with other synthetic organic polymers (polyacrylamide) for preventing seal formation, runoffs and erosion in agricultural soils exposed to heavy deluges (Tang 2006).

3.2. Carbon mineralization

In addition, the initial (C/N) ratios are 3.22 for soil and 3.25 for the (soil + PG) mixture which are favorable to the mineralisation of the organic matter in soil. The investigation of the rate of release of carbon oxide (fig. n°3) from the soil and from a (soil + PG) mixture, revealed the mineralization process of the carbon in soil. The mineralization rates are maximum at the beginning of the incubations. They are more important with (soil + PG) mixture than with soil. The rates of mineralization decrease rapidly during the incubations.

This reaction followed the first kinetic order, as described in literature (FAO). Furthermore, PG

facilitated organic matter decomposition and especially after one month of incubation at 28°C.

The fitting of this mineralization kinetic according to an exponential law ($C_t = C_0 (1 - e^{-kt})$) (Annabi 2009) led to the determination of the amount of labile compartment in soil C_0 (Saidi 2015).

C_t (mg C kg^{-1}) is the total amount of mineralized carbon at the duration of incubation t (day) per one kilogramme of soil, and k is the mineralization reaction rate constant (day^{-1}). The results are gathered in the following table 1.

Table 1: Kinetic parameters of the mineralization reaction

	Soil	PG + Soil
C (mg C kg^{-1})	378.2	399.3
k (day^{-1})	0.048	0.055

At the end of this incubation period 85% of COT in soil and 91% of COT in (soil + PG) mixture were mineralised, which is similar to the results found in cultivated soil in north Tunisia (Annabi 2009).

This result reveals that besides the utility of adding the PG to ameliorate some physicochemical properties of soil, it may alter the reserve of carbon in soil by accelerating the mineralization process. Amending PG in soils has an inhibitory effects on the reproduction of terrestrial invertebrates like earthworm known to be resources and factors for the stabilisation of organic carbon in soil (hentati earth worm). In addition, acid traces in PG contribute to the decomposition and dissolution of the organic carbon in soil, similarly to the effect of acidic rain on soil (Francis1982).

4. Conclusion

This research was performed in order to investigate the effect of PG on rain water incubated and on soil respiration. The results found can help with better understanding of PG behavior in the environment. Amended in agricultural soil, PG is an efficient fertilizer enhancing crop productivity, as it is solved by rain or irrigation water and then delivering some nutrients. Nevertheless, trace impurities in PG like acids may increase the kinetic of the mineralization of the organic carbon which has an impoverishment effect on soil. Combined amendments PG with manure or compost are planned to be studied to reveal their global effect on the soil.

Acknowledgements

Authors are grateful to IRA Medenine for financial support.

References

- Akhmetov A.S., Dmitrieva N.V. 1992. Application of phosphogypsum in road construction. Technology of mineral fertilizers, Leningrad, 19, pp. 113-115

- Annabi M., Bahri H., Latiri K. 2009. Statut organiques et respiration microbienne des sols du Nord de la Tunisie. *Biotechnol. Agron. Soc. Environ.*, 13, pp. 401-408.
- Ardhaoui K., Karbout N., Thabti S., Moussa M. 2017. *Journal of new sciences, Agriculture and Biotechnology*, CSIEA (26), 2790-2794
- Ben Amor F., Jomaa S. 2012. Regional Pilot Projects for MED POL National Action Plan, United Nations Environment Programme, GFL 4A05-2731-120600
- Bremner J.M., Mulvaney C.S. 1982. Nitrogen-total. In: Page, A.L. (Ed.), *Methods of Soil Analysis*, Part 2. American Society of Agronomy and Soil Science Society of America, Madison, pp. 595-624
- Carvalho M., Raij B. 1997. Calcium sulphate, phosphogypsum and calcium carbonate in the amelioration of acid subsoils for root growth. *Plant and Soil* 192: pp. 37-48.
- Cherifa H., Ayaria F., Ouzaria H., Marzoratib M., Brusettib L., Jedidia N., Hassena A., Daffonchio B.D. 2009. *European journal of soil biology* 45: pp. 138-145.
- Degirmenci N. 2008. The using of waste phosphogypsum and natural gypsum in adobe stabilization. *Construction and Building Materials*, 22, pp. 1220-1224.
- Drouet C. 2015. *J. Chem. Thermodynamics* 81: pp. 143-159
- Dugain F., Arial G., Audry P., Jouga J. 1961. Les méthodes d'analyses utilisées au laboratoire de physico-chimie des Sols. Dakar : ORSTOM, multigr. P.73 <http://www.documentation.ird.fr/hor/fdi:10754>
- Enamorado S., Abril J.M., Delgado A., Más J.L., Polvillo O., Quintera J.M. 2014. Implications for food safety of the uptake by tomato of 25 trace-elements from a phosphogypsum amended soil from SW Spain. *J. Hazard. Mater.* 226, pp. 122-131 <http://dx.doi.org/10.1016/j.jhazmat.2013.12.019>
- Endovitsky A.P., Batukaev A.A., Minkina T.M., Kalinitchenko V.P., Mandzhieva S.S., Sushkova S.N., Mischenko N.A., Bakoyev S. Y., Zarmaev A.A., Jusupov V.U. 2016. Carbonate Calcium Equilibrium in Soil Solution as a Driver of Heavy Metals Mobility. *Journal of Geochemical Exploration*, 2, pp. 136-153.
- FAO. 2015. Mise en valeur et amélioration des terres. Archives des documents de la FAO <http://www.fao.org/docrep/003/S8500F/s8500f0i.htm>
- Francis A.J. 1982. *Water Air Soil Pollution*. 18: 375. <https://doi.org/10.1007/BF02419425>
- Garrido F., Illera V., Campbell C.G., Garcia-Gonzalez M.T. 2006. Regulating the mobility of Cd, Cu and Pb in acid soil with amendments of phosphogypsum, sugar foam, and phosphoric rock. *European Journal of Soil Science* 57: 95-105
- Genkin M.V., Evtushenko A.V., Komkov A. A., Safiulina A. M., Spiridonov V.S., Shvetsov S.V. 2013. A method for extracting rare-earth metals and obtaining building gypsum from phosphogypsum hemihydrate. Patent RU 2187888 Russian Federation, IGC C22B 59/00, C22B 3/06.; The applicant and the patent owner United Chemical Company URALCHEM Open Joint-Stock Company – No. 2013109740/02, published on 03/03/2013, Bulletin No. 26
- Hentati O., Abrantes N., Caetano A.L., Bouguerra S., Gonçalves F., Römbke J., Pereira R. 2015. *Journal of Hazardous Materials* 294: 80-89 <https://doi.org/10.1016/j.jhazmat.2015.03.034>
- Jeddi K., Chaieb M. (2010) *Flora* 205: 184-189
- Kayouli C. 2006. Country Pasture/Forage Resource Profile Tunisia, Food and Agriculture Organization of the United Nations FAO
- Kumpiene J., Lagerkvist A., Maurice C. 2008. Stabilisation of As, Cr, Cu, Pb and Zn in soil using amendments-a review. *Waste Management*, 28, pp. 215-225
- Mackay A.D., Syers J.K. 1986. Effect of phosphate, calcium, and pH on the dissolution of a phosphate rock in soil. *Fert. Res.*, 10: 175-184.
- Mullins G.L., Mitchell C.C., Wheat forage response to tillage and sulfur applied as PG, in: *Proceedings of the third international symposium on PG*, Orlando, USA, 1990, pp. 362-375.
- Raich J.W., Rastetter E.B., Melillo J.M., Kicklighter D.W., Steudler P.A., Peterson B.J., Grace A.L., Moore B., Voëroésmarty C.J. (1991). Potential net primary productivity in South America: Application of a global model. *Ecological Applications*, 1, pp. 399-429
- Rastetter E.B., Ryan M.G., Shaver G.R., Melillo J.M., Nadelhoffer K.J., Hobbie J.E., Aber J.D. 1991. A general biogeochemistry model describing the responses of the C and N cycles in terrestrial ecosystems to changes in CO₂, climate and N deposition. *Tree Physiology*, 9: 101-126
- Robinson G.W. 1922. Particle size measurement. *J. Agr. Sci.* 12, pp. 306-321.
- Saidi N., Akef S., Bouzaiane O., Kallali H., Ben Aissa N., M'Hiri F., Jedidi N., Downer J. 2015. Carbon and nitrogen mineralization in clay loam soil amended with different composts. *Advances in Applied Agricultural Science*, 3, pp. 20-34.
- Shainberg I., Sumner M.E., Miller W.P., Farina M., Pavan M., Fey M.Y. 1989. *Use of Gypsum on Soils: A Review*. Springer-Verlag New York Inc. *Advances in Soil Science*, DOI: 10.1007/978-1-4612-3532-3_1
- Sizyakov V.M., Nutrihina S.V., Levin B.V. 2012. Technology of complex processing of phosphogypsum by conversion method to produce ammonium sulfate, phosphomel and new products. *Journal of Mining Institute*, 119, pp. 239-244.
- Tang Z., Lei T., Yu J., Shainberg I., Mamedov A., Ben-Hur M., Levy G.J., Runoff and interrill erosion in sodic soils treated with dry PAM and phosphogypsum, *Soil Sci. Soc. Am. J.*, 70, pp. 679-690.
- Tayibi H., Choura M., López F.A., Alguacil F.J., López-Delgado A. 2009. Environmental impact and management of phosphogypsum. *Journal of Environmental Management*, 90, pp. 2377-2386.
- Toma M., Saigusa M. 1997. Effects of phosphogypsum on ameliorating strongly acid nonallophanic andosols, *Plant Soil*, 192, pp. 49-55.
- Uchimiyama M., Lima I.M., Klasson K.T., Wartelle L.H. 2010. Contaminant immobilization and nutrient release by biochar soil amendment: roles of natural organic matter. *Chemosphere*, 80, pp. 935-940.
- Walkley A., black I.A. 1934. An examination method for determination for role organic matter and proposed modification of the chromic acid titration method. *Soil Sci.*, 37, pp. 29-38.
- Wright R.J., Kemper W.D., Millner P.D., Power J.F., and Korkak R. F. 1998. *Agricultural Uses of Municipal, Animal, and Industrial Byproducts* U.S. Department of Agriculture, Agricultural Research Service, Conservation Research Report 44 Chapter 7 pp 120
- Wu H., Guo Z., Peng C. 2003. Land use induced changes of organic carbon storage in soils of China. *Global Change Biology*, 9, pp. 305-315.
- You H., Jin H., Khaldi A., Kwak M., Lee T., Khaine I., Jang J., Lee H., Kim I., Ahn T., Song J., Song Y., Khorchani A., Stiti B., Woo S. 2016. Plant diversity in different bioclimatic zones in Tunisia. *Journal of Asia-Pacific Biodiversity*, 9, pp. 56-62.
- Zairi M., Rouis M.J. 1999. Impacts environnementaux du stockage du phosphogypse à Sfax (Tunisie). *Bulletin Des Laboratoires Des Ponts Et Chaussées*, 219, pp. 29-40.
- Zoghalmi R.I., Ben aissa N., Hamdi H., Mokni-Tlili S., Khelil M.N., Jedidi N. 2016. Minéralisation du carbone organiques et sa corrélation avec la biomasse microbienne dans deux sols agricoles suite à un amendement de boues résiduaires. *Journal of New Sciences, Agriculture and Biotechnology*, 31, pp. 1812-1821